

11) Publication number:

0 347 146 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (4) Date of publication of patent specification: 01.09.93 (5) Int. Cl.⁵: C07D 471/04, C07D 487/04, C07D 475/02, A61K 31/505,
- 21 Application number: 89305911.3
- 2 Date of filing: 12.06.89

C07D 471/04, C07D 487/04, C07D 475/02, A61K 31/505, A61K 31/53, C07D 213/81, C07D 241/24, C07D 253/06, //(C07D471/04,239:00,221:00), (C07D487/04,241:00,239:00), (C07D487/04,253:00,239:00)

- So Fused pyrimidine derivatives, process and intermediates for their preparation and pharmaceutical compositions containing them.
- Priority: 16.06.88 GB 8814351 16.06.88 GB 8814350 16.06.88 GB 8814353
- ② Date of publication of application: 20.12.89 Bulletin 89/51
- Publication of the grant of the patent: 01.09.93 Bulletin 93/35
- Designated Contracting States:
 AT BE CH DE ES FR GB GR IT LI LU NL SE
- 99 References cited: DE-A- 1 795 722 FR-A- 2 225 166 GB-A- 1 543 874
- Proprietor: SMITH KLINE & FRENCH LABORA-TORIES LIMITED

Mundells Welwyn Garden City Hertfordshire, AL7 1EY(GB)

- ② Inventor: Coates, William John
 SmithKline Pharmaceuticals, The Frythe,
 Welwyn, Hertfordshire AL6 9AR(GB)
 Inventor: Flynn, Sean Thomas
 SmithKline Pharmaceuticals, The Frythe,
 Welwyn, Hertfordshire AL6 9AR(GB)
 Inventor: Rawlings, Derek Anthony
 SmithKline Pharmaceuticals, The Frythe,
 Welwyn, Hertfordshire AL6 9AR(GB)
- (2) Representative: Waters, David Martin, Dr. et al Smith Kline & French Laboratories Ltd. Patent Department Mundells Welwyn Garden City Hertfordshire AL7 1EY (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Descripti n

20

25

30

50

55

The present invention relates to fused pyrimidine derivatives, processes for their preparation, intermediates in their preparation, their use as therapeutic agents and to pharmaceutical compositions containing them. The compounds of this invention are inhibitors of a calmodulin insensitive cyclic GMP phosphodiesterase and are of use in combating such conditions where such inhibition is thought to be beneficial. They are bronchodilators and are therefore of use in combating chronic reversible obstructive lung diseases such as asthma and bronchitis. Some of the compounds of the present invention have antiallergic activity and are therefore useful in combating allergic diseases such as allergic asthma, allergic rhinitis, urticaria and irritable bowel syndrome. Furthermore the compounds of this invention are vasodilators and are therefore of value in combating angina, hypertension and congestive heart failure.

GB-A-1,543,874 (Carlo Erba) discloses a series of substituted phenyl-3,4-dihydro-4-oxo-quinazoline derivatives which are said to possess anti-allergic activity.

FR-A-2,225,166 (Pfizer) discloses bicyclic and tricyclic ring systems containing a pyrimidine moiety as anti-allergic agents.


DE-A-1,795,722 (Karl Thomae) discloses substituted pyrimido[5,4-d]pyrimidines which act as cardiovascular agents.

Accordingly, the present invention provides compounds of the formula (1):

(1)

and pharmaceutically acceptable salts thereof, wherein

is a ring of sub-formula (a), (b), (c), (d), (e), (f) or (g):



R1 is C₁₋₆ alkyl, C₂₋₆ alkenyl, C₃₋₅ cycloalkylC₁₋₆ alkyl, or C₁₋₆ alkyl substituted by 1 to 6 fluoro groups;

R² is C_{1-6} alkylthio, C_{1-6} alkylsulphonyl, C_{1-6} alkoxy, hydroxy, hydrogen, hydrazino, C_{1-6} alkyl, phenyl, -NHCOR³ wherein R³ is hydrogen or C_{1-6} alkyl, or -NR⁴R⁵, wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, hexahydroazepino, morpholino or piperazino ring, or R⁴ and R⁵ are independently hydrogen, C_{3-5} cycloalkyl or C_{1-6} alkyl which is optionally substituted by -CF₃, phenyl, -S(O)_nC₁₋₆ alkyl wherein n is 0, 1 or 2, -OR⁶, -CO₂R⁷ or -NR⁶R՞ wherein R⁶ to R³ are independently hydrogen or C_{1-6} alkyl, provided that the carbon atom adjacent to the nitrogen atom is not substituted by said -S(O)_nC₁₋₆ alkyl, -OR⁶ or -NR⁶R³ groups; and

R is hydrogen and can also be hydroxy when R2 is hydroxy.

Suitably R1 is C2-5 alkyl for example ethyl, n-propyl, isopropyl, butyl, isobutyl or pentyl.

Suitably R^1 is C_{3-5} alkenyl for example propenyl, butenyl or pentenyl.

Suitably R1 is cyclopropylmethyl.

Examples of C₁₋₆ alkyl substituted by 1 to 6 fluoro groups include -CF₃, -CH₂CF₃ or -CF₂CHFCF₃.

Preferably R1 is n-propyl.

Suitably R² is C₁₋₆ alkylthio, C₁₋₆ alkylsulphonyl or C₁₋₆ alkoxy for example methylthio, ethylthio, methylsulphonyl, ethylsulphonyl, methoxy, ethoxy or propoxy.

Suitably R² is hydroxy, hydrogen or hydrazino.

Suitably R^2 is phenyl or C_{1-6} alkyl for example methyl, ethyl or propyl.

Suitably R² is -NHCOR³ for example formamido or acetamido.

Suitably R² is -NR⁴R⁵ for example amino, methylamino, ethylamino, propylamino, dimethylamino, diethylamino, dipropylamino, cyclopropylamino, morpholino, 2,2,2-trifluoroethylamino, phenethylamino, 3-methylthiopropylamino, 3-methylsulphinylpropylamino, 3-methylsulphonylpropylamino, 2-hydroxyethylamino, 3-methoxypropylamino, N-ethyl-N-(2-hydroxyethyl)amino, 2-aminoethylamino, 2-dimethylaminoethylamino, ethoxycarbonylmethylamino, carboxymethylamino, 2-ethoxycarbonylethylamino or 2-carboxyethylamino.

roonylethylamino or z-carboxyethylamino

Suitably

55

25

A

is a group of sub-formula (a) thus forming a pyrido[2,3-d]pyrimidine ring system.

Suitably



is a group of sub-formula (b) thus forming a pyrido[3,4-d]pyrimidine ring system. Suitably

10

15



is a group of sub-formula (c) thus forming a pyrido[4,3-d]pyrimidine ring system. Suitably



20

25

30

is a group of sub-formula (d) thus forming a pyrido[3,2-d]pyrimidine ring system. Suitably

A

is a group of sub-formula (e) thus forming a pteridine ring system. Suitably

YA

35 is a group of sub-formula (f) thus forming a pyrimido[5,4-e][1,2,4]triazine ring system. Suitably



40

is a group of sub-formula (g) thus forming a pyrimido[4,5-e][1,2,4]triazine ring system. Particular compounds of this invention are:

2-(2-propoxyphenyl)pyrido[2,3-d]pyrimid-4(3H)-one,

2-(2-propoxyphenyl)pyrido[3,4-d]pyrimid-4(3H)-one,

2-(2-propoxyphenyl)pyrido[4,3-d]pyrimid-4(3H)-one,

2-(2-propoxyphenyl)pyrido[3,2-d]pyrimid-4(3H)-one,

2-(2-propoxyphenyl)pteridin-4(3H)-one,

2-(2-propoxyphenyl)pteridin-4,6(3H,5H)-dione,

2-(2-propoxyphenyl)pteridin-4,6,7(3H,5H,8H)-trione,

5,6-dihydro-3-methylthio-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,

3-amino-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,

3-methylamino-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,

3-methoxy-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,

3-methylthio-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,

3-amino-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,

3-methylamino-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,

3-methoxy-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,